

Immunization 101



Katie Reilly, MPH, MSN, RN, APHN-BC
Massachusetts Department of Public Health
Immunization Program

Presenter Disclosure Information

I, Katie Reilly, have been asked to disclose any significant relationships with commercial entities that are either providing financial support for this program or whose products or services are mentioned during my presentations. I have no relationships to disclose.

I may discuss the use of vaccines in a manner not approved by the U.S. Food and Drug Administration

- But in accordance with ACIP recommendations

Outline

- Herd Immunity
- Types of vaccines
- 2017 Adult Immunization Schedule
- Screening prior to vaccination
- Contraindications and precautions to vaccination
- Vaccine Safety
- Vaccine Information Statements (VIS)
- Vaccine administration documentation requirements
- Vaccine adverse events and medical error reporting
- Use of Model Standing Orders

Herd Immunity/Community Immunity

“A situation in which a sufficient proportion of a population is immune to an infectious disease (through vaccination and/or prior illness) to make its spread from person to person unlikely. Even individuals not vaccinated (such as newborns and those with chronic illnesses) are offered some protection because the disease has little opportunity to spread within the community.”

Retrieved from:

<http://www.cdc.gov/vaccines/terms/glossary.html#communityimmunity>




Photo credit: Courtesy: The National Institute of Allergy and Infectious Disease (NIAID)

Live Attenuated Vaccines

- Attenuated (weakened) form of the “wild” virus or bacterium
- Must replicate to produce an immune response
- Immune response virtually identical to natural infection
- Usually produce immunity with one dose (except those administered orally)
- Interference from circulating antibody
- Fragile: must be stored and handled carefully
- Viral: measles, mumps, rubella, vaccinia, varicella, zoster, yellow fever, rotavirus, intranasal influenza, oral polio*
- Bacterial: BCG*, oral typhoid

*not available in the USA

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/prinvac.pdf>

Inactivated Vaccines

- Cannot replicate, and therefore cannot cause infection
- Less affected by circulating antibody than live vaccine
- Always require multiple doses
- Immune response mostly humoral
- Antibody titer diminish with time
- May require periodic supplemental booster doses
- Whole cell vaccines:
 - Viral: polio, hepatitis A, rabies, influenza*
 - Bacterial: pertussis*, typhoid*, cholera, plague*
- Fractional vaccines
 - Subunits: hepatitis B, influenza, acellular pertussis, HPV, anthrax
 - Toxoids: diphtheria, tetanus

*not available in the USA

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/prinvac.pdf>

2017 Adult Immunization Schedule



MMWR 2017:66:136.

Annals of Internal Medicine 2017;166:209.



Available at:

- <https://www.cdc.gov/mmwr/volumes/66/wr/pdfs/mm6605e2.pdf>
- <https://www.cdc.gov/vaccines/schedules/>
- <http://annals.org/aim/article/2601209/recommended-immunization-schedule-adults-aged-19-years-older-united-states>

Updates - 2017 Adult Immunization Schedule

- **Influenza vaccination**
 - LAIV not recommended in 2016–2017 season
 - Modified language on egg allergy
- **Tdap vaccination**
 - Guidance for use during pregnancy
- **HPV vaccination**
 - Updated dosing schedule
- **Hepatitis B vaccination**
 - Updated definition of chronic liver disease
- **Meningococcal vaccination**
 - Use of MenACWY for adults with HIV infection
 - Updated dosing schedule for MenB-FHbp
- **Format changes**
 - New first page of resources, references and abbreviations
 - Incorporated human factors and ergonomics assessment
 - Modified footnotes for consistency
 - Integrated table of contraindications and precautions

Figures 1 and 2 should be read with the footnotes that contain information about special populations.

Age Groups 27-49 & 50-59 Combined

Figure 1. Recommended immunization schedule for adults aged 19 years or older by age group, unless stated otherwise.

Vaccine	19–21 years	22–26 years	27–59 years	60–64 years	≥ 65 years
Influenza ¹	1 dose annually				
Td/Tdap ²	Substitute Tdap for Td once, then Td booster every 10 yrs				
MMR ³	1 or 2 doses depending on indication				
VAR ⁴	2 doses				
HZV ⁵				1 dose	
HPV–Female ⁶	3 doses				
HPV–Male ⁶	3 doses				
PCV13 ⁷					1 dose
PPSV23 ⁷	1 or 2 doses depending on indication				1 dose
HepA ⁸	2 or 3 doses depending on vaccine				
HepB ⁹	3 doses				
MenACWY or MPSV4 ¹⁰	1 or more doses depending on indication				
MenB ¹⁰	2 or 3 doses depending on vaccine				
Hib ¹¹	1 or 3 doses depending on indication				

Live Vaccines Grouped

 Recommended for adults who meet the age requirement, lack documentation of vaccination, or lack evidence of past infection
  Recommended for adults with additional medical conditions or other indications
  No recommendation

Figure 2. Recommended immunization schedule for adults aged 19 years or older by medical condition and other indications, United States, 2017

Vaccine	Pregnancy ^{1-6,9}	Immuno-compromised (excluding HIV infection) ^{3-7,11}	HIV infection CD4+ count (cells/ μ L) ^{3-7,9-11}		Asplenia, persistent complement deficiencies ^{7,10,11}	Kidney failure, end-stage renal disease, on hemodialysis ^{7,9}	Heart or lung disease, chronic alcoholism ⁷	Chronic liver disease ⁷⁻⁹	Diabetes ^{7,9}	Healthcare personnel ^{3,4,9}	Men who have sex with men ^{6,8,9}
			< 200	\geq 200							
Influenza ¹	1 dose annually										
Td/Tdap ²	1 dose Tdap each pregnancy	Substitute Tdap for Td once, then Td booster every 10 yrs									
MMR ³	contraindicated		1 or 2 doses depending on indication								
VAR ⁴	contraindicated		2 doses								
HZV ⁵	contraindicated			1 dose							
HPV-Female ⁶		3 doses through age 26 yrs									
HPV-Male ⁶		3 doses through age 26 yrs			3 doses through age 21 yrs						3 doses through age 26 yrs
PCV13 ⁷		1 dose									
PPSV23 ⁷		1, 2, or 3 doses depending on indication									
HepA ⁸	2 or 3 doses depending on vaccine										
HepB ⁹	3 doses										
MenACWY or MPSV4 ¹⁰	HIV Added		1 or more doses depending on indication								
MenB ¹⁰	2 or 3 doses depending on vaccine										
Hib ¹¹		3 doses post-HSCT recipients only	1 dose								



Recommended for adults who meet the age requirement, lack documentation of vaccination, or lack evidence of past infection



Recommended for adults with additional medical conditions or other indications



Contraindicated



No recommendation

Screening

- Is key to preventing serious adverse reactions
- Specific questions intended to identify contraindications or precautions to vaccination
- Screening must occur at every immunization encounter (not just before the first dose)
- Use of a standardized form will facilitate effective screening

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/genrec.pdf>

Immunization Action Coalition (IAC) Screening Forms

- Child and Teen Immunizations
- Adult Immunizations
- IIV Seasonal Influenza

<http://www.immunize.org/handouts/screening-vaccines.asp>

Screening Checklist for Contraindications to Vaccines for Adults

PATIENT NAME _____

DATE OF BIRTH _____ / _____ / _____
month / day / year

For patients: The following questions will help us determine which vaccines you may be given today. If you answer "yes" to any question, it does not necessarily mean you should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

	yes	no	don't know
1. Are you sick today?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Do you have allergies to medications, food, a vaccine component, or latex?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Have you ever had a serious reaction after receiving a vaccination?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Do you have a long-term health problem with heart disease, lung disease, asthma, kidney disease, metabolic disease (e.g., diabetes), anemia, or other blood disorder?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Do you have cancer, leukemia, HIV/AIDS, or any other immune system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Information for Healthcare Professionals about the Screening Checklist for Contraindications to Vaccines for Adults

Are you interested in knowing why we included a certain question on the screening checklist? If so, read the information below. If you want to find out even more, consult the references listed at the end.

- 1. Are you sick today?** [all vaccines]
There is no evidence that acute illness reduces vaccine efficacy or increases vaccine adverse events.¹ However, as a precaution with moderate or severe acute illness, all vaccines should be delayed until the illness has improved. Mild illnesses (such as upper respiratory infections or diarrhea) are NOT contraindications to vaccination. Do not withhold vaccination if a person is taking antibiotics.
- 2. Do you have allergies to medications, food, a vaccine component, or latex?** [all vaccines]
An anaphylactic reaction to latex is a contraindication to vaccines that contain latex as a component or as part of the packaging (e.g., vial stoppers, prefilled syringe plungers, prefilled syringe caps). If a person has anaphylaxis after eating gelatin, do not administer vaccines containing gelatin. A local reaction to a prior vaccine dose or vaccine component, including latex, is not a contraindication to a subsequent dose or vaccine containing that component. For information on vaccines supplied in vials or syringes containing latex, see reference 2; for an extensive list of vaccine components, see reference 3.
People with egg allergy of any severity can receive any recommended influenza vaccine (i.e., any IIV or RIV) that is otherwise appropriate for the patient's age. For people with a history of severe allergic reaction to egg involving any symptom other than hives (e.g., angioedema, respiratory distress), or who required epinephrine or another emergency medical intervention, the vaccine should be administered in a medical setting, such as a clinic, health department, or physician office. Vaccine administration should be supervised by a healthcare provider who is able to recognize and manage severe allergic conditions.⁴
- 3. Have you ever had a serious reaction after receiving a vaccination?** [all vaccines]
History of anaphylactic reaction (see question 2) to a previous dose of vaccine or vaccine component is a contraindication for subsequent doses.¹ Under normal circumstances, vaccines are deferred when a precaution is present. However, situations may arise when the benefit outweighs the risk (e.g., during a community pertussis outbreak).
- 4. Do you have a long-term health problem with heart disease, lung disease, asthma, kidney disease, metabolic disease (e.g., diabetes), anemia, or other blood disorder?** [LAIV]
The safety of intranasal live attenuated influenza vaccine (LAIV) in people with these conditions has not been established. These conditions, including asthma in

NOTE: Live attenuated influenza vaccine (LAIV4; FluMist), is not recommended by CDC's Advisory Committee on Immunization Practices for use in the U.S. during the 2016–17 influenza season. Because LAIV4 is still a licensed vaccine that might be available and that some providers might elect to use, for informational purposes, reference is made to previous recommendations for its use.

- 7. Have you had a seizure or a brain or other nervous system problem?** [influenza, Td/Tdap]
Tdap is contraindicated in people who have a history of encephalopathy within 7 days following DTP/DTaP given before age 7 years. An unstable progressive neurologic problem is a precaution to the use of Tdap. For people with stable neurologic disorders (including seizures) unrelated to vaccination, or for people with a family history of seizure, vaccinate as usual. A history of Guillain-Barré syndrome (GBS) is a consideration with the following: 1) Td/Tdap: if GBS has occurred within 6 weeks of a tetanus-containing vaccine and decision is made to continue vaccination, give Tdap instead of Td if no history of prior Tdap; 2) Influenza vaccine (IIV/LAIV): if GBS has occurred within 6 weeks of a prior influenza vaccine, vaccinate with IIV if at increased risk for severe influenza complications.
- 8. During the past year, have you received a transfusion of blood or blood products, or been given immune (gamma) globulin or an antiviral drug?** [LAIV, MMR, VAR, ZOS]
Certain live virus vaccines (e.g., LAIV, MMR, VAR, ZOS) may need to be deferred, depending on several variables. Consult the most current ACIP recommendations for current information on intervals between antiviral drugs, immune globulin or blood product administration and live virus vaccines.¹
- 9. For women: Are you pregnant or is there a chance you could become pregnant during the next month?** [HPV, IIV, MMR, LAIV, VAR, ZOS]
Live virus vaccines (e.g., MMR, VAR, ZOS, LAIV) are contraindicated one month before and during pregnancy because of the theoretical risk of virus transmission to the fetus. Sexually active women in their childbearing years who receive live virus vaccines should be instructed to practice careful contraception for one month following receipt of the vaccine. On theoretical grounds, inactivated poliovirus vaccine should not be given during pregnancy; however, it may be given if risk of exposure is imminent and immediate protection is needed (e.g., travel to endemic areas). Inactivated influenza vaccine and Tdap are both recommended during pregnancy. Both

-
-
-
-

Contraindication and Precautions

Contraindication

- A condition that increases the likelihood of a serious adverse reaction to a vaccine for a patient with that condition.
- In general, vaccine should not be administered when a contraindication condition is present.

Precaution

- A condition in a recipient that might increase the chance or severity of a serious adverse reaction, or that might compromise the ability of the vaccine to produce immunity.
- In general, vaccines are deferred with a precaution condition is present. However, situations may arise when the benefit of the protection from the vaccine outweighs the risk of an adverse reaction, and the provider may decide to give the vaccine.

Contraindications & Precautions

Summary Table Published Annually by CDC with Adult Schedule

- Adult Summary Table:

<https://www.cdc.gov/vaccines/schedules/downloads/adult/adult-combined-schedule.pdf#page=6>

Table. Contraindications and precautions for vaccines recommended for adults aged 19 years or older*

The Advisory Committee on Immunization Practices (ACIP) recommendations and package inserts for vaccines provide information on contraindications and precautions related to vaccines. Contraindications are conditions that increase chances of a serious adverse reaction in vaccine recipients and the vaccine should not be administered when a contraindication is present. Precautions should be reviewed for potential risks and benefits for vaccine recipient. For a person with a severe allergy to latex, e.g., anaphylaxis, vaccines supplied in vials or syringes that contain natural rubber latex should not be administered unless the benefit of vaccination clearly outweighs the risk for a potential allergic reaction. For latex allergies other than anaphylaxis, vaccines supplied in vials or syringes that contain dry, natural rubber or natural rubber latex may be administered.

Contraindications and precautions for vaccines routinely recommended for adults

Vaccine	Contraindications	Precautions
All vaccines routinely recommended for adults	• Severe reaction, e.g., anaphylaxis, after a previous dose or to a vaccine component	• Moderate or severe acute illness with or without fever

Additional contraindications and precautions for vaccines routinely recommended for adults

Vaccine	Additional Contraindications	Additional Precautions
IIV ¹		<ul style="list-style-type: none"> • History of Guillain-Barré Syndrome within 6 weeks after previous influenza vaccination • Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, or recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting and under the supervision of a healthcare provider who is able to recognize and manage severe allergic conditions)
RIV ¹		<ul style="list-style-type: none"> • History of Guillain-Barré Syndrome within 6 weeks after previous influenza vaccination
LAIV ¹	• LAIV should not be used during 2016–2017 influenza season	• LAIV should not be used during 2016–2017 influenza season
Tdap/Td	<ul style="list-style-type: none"> • For pertussis-containing vaccines: encephalopathy, e.g., coma, decreased level of consciousness, or prolonged seizures, not attributable to another identifiable cause within 7 days of administration of a previous dose of a vaccine containing tetanus or diphtheria toxoid or acellular pertussis 	<ul style="list-style-type: none"> • Guillain-Barré Syndrome within 6 weeks after a previous dose of tetanus toxoid-containing vaccine • History of Arthus-type hypersensitivity reactions after a previous dose of tetanus or diphtheria toxoid-containing vaccine. Defer vaccination until at least 10 years have elapsed since the last tetanus toxoid-containing vaccine • For pertussis-containing vaccine, progressive or unstable neurologic disorder, uncontrolled seizures, or progressive encephalopathy (until a treatment regimen has been established and the condition has stabilized)
MMR ²	<ul style="list-style-type: none"> • Severe immunodeficiency, e.g., hematologic and solid tumors, chemotherapy, congenital immunodeficiency or long-term immunosuppressive therapy³, human immunodeficiency virus (HIV) infection with severe immunocompromise • Pregnancy 	<ul style="list-style-type: none"> • Recent (within 11 months) receipt of antibody-containing blood product (specific interval depends on product)⁴ • History of thrombocytopenia or thrombocytopenic purpura • Need for tuberculin skin testing⁵
VAR ²	<ul style="list-style-type: none"> • Severe immunodeficiency, e.g., hematologic and solid tumors, chemotherapy, congenital immunodeficiency or long-term immunosuppressive therapy³, HIV infection with severe immunocompromise • Pregnancy 	<ul style="list-style-type: none"> • Recent (within 11 months) receipt of antibody-containing blood product (specific interval depends on product)⁴ • Receipt of specific antiviral drugs (acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination (avoid use of these antiviral drugs for 14 days after vaccination)
HZV ²	<ul style="list-style-type: none"> • Severe immunodeficiency, e.g., hematologic and solid tumors, chemotherapy, congenital immunodeficiency or long-term immunosuppressive therapy³, HIV infection with severe immunocompromise • Pregnancy 	<ul style="list-style-type: none"> • Receipt of specific antiviral drugs (acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination (avoid use of these antiviral drugs for 14 days after vaccination)
HPV vaccine		• Pregnancy


Vaccination of Pregnant Women

- Live vaccines should not be administered to women known to be pregnant
- In general, inactivated vaccines may be administered to pregnant women for whom they are indicated
- HPV vaccine should be deferred during pregnancy

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/genrec.pdf>

CDC Guidelines for Vaccinating Pregnant Women

- Guidelines for vaccination
- Travel and other vaccines
- Breastfeeding and vaccination
- Prenatal screening



<http://www.cdc.gov/vaccines/pregnancy/hcp/guidelines.html>

Immunizations and Pregnancy



Vaccine	Before pregnancy	During pregnancy	After pregnancy	Type of vaccine
Influenza	Yes	Yes, during flu season	Yes	Inactivated
Tdap	May be recommended; it is better to vaccinate during pregnancy when possible	Yes, during each pregnancy	Yes, immediately postpartum, if Tdap never received in lifetime; it is better to vaccinate during pregnancy	Toxoid/ Inactivated
Td	May be recommended	May be recommended, but Tdap is preferred	May be recommended	Toxoid
Hepatitis A	May be recommended	May be recommended	May be recommended	Inactivated
Hepatitis B	May be recommended	May be recommended	May be recommended	Inactivated
Meningococcal	May be recommended	Base decision on risk vs. benefit; inadequate data for specific recommendation	May be recommended	Inactivated
Pneumococcal	May be recommended	Base decision on risk vs. benefit; inadequate data for specific recommendation	May be recommended	Inactivated
HPV	May be recommended (through 26 years of age)	No	May be recommended (through 26 years of age)	Inactivated
MMR	May be recommended; once received, avoid conception for 4 weeks	No	May be recommended	Live
Varicella	May be recommended; once received, avoid conception for 4 weeks	No	May be recommended	Live

<http://www.cdc.gov/vaccines/pregnancy/downloads/immunizations-preg-chart.pdf>

Importance of Vaccine Safety

Decreases in disease risks and increased attention on vaccine risks

Public confidence in vaccine safety critical

- Higher standard of safety is expected of vaccines
- Vaccinees generally healthy (vs. ill for drugs)
- Lower risk tolerance = need to search for rare reactions
- Vaccination universally recommended and mandated

<http://www.cdc.gov/vaccines/pubs/pinkbook/safety.html>

US Post-licensure Vaccine Safety System

System	Collaboration	Description
Vaccine Adverse Event Reporting System (VAERS)	CDC and FDA	Frontline spontaneous reporting system to detect potential vaccine safety issues
Vaccine Safety Datalink (VSD)	CDC and 9 Integrated Health Care Systems	Large linked database system used for active surveillance and research ~9.4 million members (~3% of US pop.) -Conducts monitoring & evaluation
Clinical Immunization Safety Assessment (CISA) Project	CDC and 7 Academic Centers	Expert collaboration that conducts individual clinical vaccine safety assessments and clinical research
Post-Licensure Rapid Immunization Safety Monitoring Program (PRISM)	FDA and 4 partner organizations	Large distributed database system used for active surveillance and research ~170 million individuals

Source: HPV Safety Presentation by Julianne Gee, MPH Immunization Safety Office
Centers for Disease Control and Prevention (CDC) August 4, 2016

The Provider's Role

Immunization providers can help to ensure the safety and efficacy of vaccines through proper:

- Vaccine storage and administration
- Timing and spacing of vaccine doses
- Observation of contraindications and precautions
- Management of adverse reactions
- Reporting to VAERS
- Benefit and risk communication

<http://www.cdc.gov/vaccines/pubs/pinkbook/safety.html>

Seven Rights of Vaccine Administration

- Right Patient
- Right Time
- Right Vaccine (and Diluent)
- Right Dosage
- Right Route, Needle, Technique
- Right Injection Site
- Right Documentation

<http://www.immunize.org/technically-speaking/20141101.asp>

Influenza Vaccine Products for the 2016–2017 Influenza Season

Manufacturer	Trade Name (vaccine abbreviation) ¹	How Supplied	Mercury Content (µg Hg/0.5mL)	Age Group	Vaccine Product Billing Code ²	
					CPT	Medicare ³
AstraZeneca	FluMist ⁴ (LAIV4)	0.2 mL (single-use nasal spray)	0	2 through 49 years	90672	90672
GlaxoSmithKline	Fluarix (IIV4)	0.5 mL (single-dose syringe)	0	3 years & older	90686	90686
ID Biomedical Corp. of Quebec, a subsidiary of GlaxoSmithKline	FluLaval (IIV4)	0.5 mL (single-dose syringe)	0	6 months & older	90686	90686
		5.0 mL (multi-dose vial)	<25	6 months & older	90688	90688
Protein Sciences Corp.	Flublok (RIV3)	0.5 mL (single-dose vial)	0	18 years & older	90673	90673
Sanofi Pasteur, Inc.	Fluzone (IIV4)	0.25 mL (single-dose syringe)	0	6 through 35 months	90685	90685
		0.5 mL (single-dose syringe)	0	3 years & older	90686	90686
		0.5 mL (single-dose vial)	0	3 years & older	90686	90686
		5.0 mL (multi-dose vial)	25	6 through 35 months	90687	90687
		5.0 mL (multi-dose vial)	25	3 years & older	90688	90688
	Fluzone High-Dose (IIV3-HD)	0.5 mL (single-dose syringe)	0	65 years & older	90662	90662
	Fluzone Intradermal (IIV4-ID)	0.1 mL (single-dose microinjection system)	0	18 through 64 years	90630	90630
Seqirus (formerly Novartis influenza vaccines and bioCSL)	Afluria (IIV3)	0.5 mL (single-dose syringe)	0	9 years & older ^{5,6}	90656	90656
		5.0 mL (multi-dose vial)	24.5		90658	Q2035
	Afluria (IIV4)	0.5 mL (single-dose syringe)	0	18 years & older ⁶	90686	90686
		5.0 mL (multi-dose vial)	24.5		90688	90688
	Fluad (aIIV3)	0.5 mL (single-dose syringe)	0	65 years & older	90653	90653
	Fluvirin (IIV3)	0.5 mL (single-dose syringe)	≤1	4 years & older	90656	90656
		5.0 mL (multi-dose vial)	25		90658	Q2037
	Flucelvax (ccIIV4)	0.5 mL (single-dose syringe)	0	4 years & older	90674	90674

<http://www.immunize.org/catg.d/p4072.pdf>

Administering Vaccines to Adults: Dose, Route, Site, and Needle Size

VACCINE	DOSE	ROUTE
Hepatitis A (HepA)	≤18 yrs: 0.5 mL ≥19 yrs: 1.0 mL	
Hepatitis B (HepB)	≤19 yrs: 0.5 mL ≥20 yrs: 1.0 mL	
HepA-HepB (Twinrix)	≥18 yrs: 1.0 mL	
Human papillomavirus (HPV)	0.5 mL	
Influenza, live attenuated (LAIV)	0.2 mL (0.1 mL into each nostril)	NAS
Influenza, inactivated (IIV) and recombinant (RIV)	0.5 mL	
Influenza (IIV) Fluzone Intradermal, for ages 18 through 64 years	0.1 mL	ID
Measles, Mumps, Rubella (MMR)	0.5 mL	
Meningococcal conjugate (MenACWY)	0.5 mL	
Meningococcal protein (MenB)	0.5 mL	
Meningococcal serogroup B (MenB)	0.5 mL	
Meningococcal polysaccharide (MPSV)	0.5 mL	
Pneumococcal conjugate (PCV13)	0.5 mL	
Pneumococcal polysaccharide (PPSV)	0.5 mL	I
Tetanus, Diphtheria (Td) with Pertussis (Tdap)	0.5 mL	
Varicella (VAR)	0.5 mL	

Intramuscular (IM) injection

Vaccines with Diluents: How to Use Them

Be sure to reconstitute the following vaccines correctly before administering them! Reconstitution means that the lyophilized (freeze-dried) vaccine powder or wafer in one vial must be reconstituted (mixed) with the diluent (liquid) in another.

- Only use the diluent provided by the manufacturer for that vaccine as indicated on the chart.
- ALWAYS check the expiration date on the diluent and vaccine. NEVER use expired diluent or vaccine.

Vaccine product name	Manufacturer	Lyophilized vaccine (powder)	Liquid diluent (may contain vaccine)	Time allowed between reconstitution and use, as stated in package insert ^a	Diluent storage environment
ActHIB (Hib)	Sanofi Pasteur	Hib	0.4% sodium chloride	24 hrs	Refrigerator
Hiberix (Hib)	GlaxoSmithKline	Hib	0.9% sodium chloride	24 hrs	Refrigerator or room temp
Imovax (RAB _{HDCV})	Sanofi Pasteur	Rabies virus	Sterile water	Immediately [†]	Refrigerator
M-M-R II (MMR)	Merck	MMR	Sterile water	8 hrs	Refrigerator or room temp
MenHibrix (Hib-MenCY)	GlaxoSmithKline	Hib-MenCY	0.9% sodium chloride	Immediately [†]	Refrigerator or room temp
Menomune (MPSV4)	Sanofi Pasteur	MPSV4	Distilled water	Single-dose vial: Immediately [†] Multidose vial: 35 days	Refrigerator
Menveo (MenACWY)	GlaxoSmithKline	MenA	MenCWY	8 hrs	Refrigerator
Pentacel (DTaP-IPV/Hib)	Sanofi Pasteur	Hib	DTaP-IPV	Immediately [†]	Refrigerator
ProQuad (MMRV)	Merck	MMRV	Sterile water	30 min	Refrigerator or room temp
RabAvert (RAB _{PECEV})	GlaxoSmithKline	Rabies virus	Sterile water	Immediately [†]	Refrigerator
Rotarix (RV1) [‡]	GlaxoSmithKline	RV1	Sterile water, calcium carbonate, and xanthan	24 hrs	Refrigerator or room temp
Varivax (VAR)	Merck	VAR	Sterile water	30 min	Refrigerator or room temp
YF-VAX (YF)	Sanofi Pasteur	YF	0.9% sodium chloride	60 min	Refrigerator
Zostavax (HZV)	Merck	HZV	Sterile water	30 min	Refrigerator or room temp

<http://www.immunize.org/catg.d/p3084.pdf>


<http://www.immunize.org/catg.d/p3040.pdf>

One & Only Campaign: Injection Safety Guidelines

- Follow proper infection control practices and maintain aseptic technique during the preparation and administration of injected medications
- Never administer medications from the same syringe to more than one patient, even if the needle is changed
- Never enter a vial with a used syringe or needle
- Do not use medications packaged as a single-dose or single-use for more than one patient
- Whenever possible and appropriate, limit use of multi-dose vials

CDC & Safe Injection Practices Coalition
<http://www.cdc.gov/injectionsafety/1anOnly.html>

MDPH Adult Immunization Conference 2017



Vaccine Information Statements (VISs)


Healthcare provider requirements

- Public and private providers
- Give VISs **before** vaccine is administered
- Applies to **every dose** of a vaccine series not just the first dose
- Opportunities for questions should be provided before each vaccination
- Offer a copy of the VISs to take away
- Available in multiple languages

<http://www.cdc.gov/vaccines/hcp/vis/about/facts-vis.html#give>

Your Sources for VISs

<http://www.cdc.gov/vaccines/hcp/vis/index.html>

 **Get Email Updates**

To receive email updates about this page, enter your email address:

[What's this?](#)


<http://www.immunize.org/vis/>

NEW New and Revised VISs
Check here for weekly updates

Current VIS Dates

Check your stock of VISs against this list. If you have outdated VISs, get current versions.

Adenovirus	6/11/14	MMR	4/20/12
Anthrax	3/10/10	MMRV	5/21/10
Chickenpox	3/13/08	Multi-vaccine	11/5/15
DTaP	5/17/07	PCV13	11/5/15
Hib	4/2/15	PPSV	4/24/15
Hepatitis A	7/20/16	Polio	7/20/16
Hepatitis B	7/20/16	Rabies	10/6/09
HPV-Cervarix	5/3/11	Rotavirus	4/15/15
HPV-Gardasil	5/17/13	Shingles	10/6/09
HPV-Gardasil 9	3/31/16	Td	2/24/15
Influenza	8/7/15	Tdap	2/24/15
J. enceph.	1/24/14	Typhoid	5/29/12
MCV4/MPSV4	3/31/16	Y. fever	3/30/11
MenB	8/9/16		

[PRINT VERSION](#) 

Healthcare Provider Documentation Requirements

Providers must ensure that the recipient's permanent medical record (whether paper-based or electronic) contains all of the required vaccine administration documentation, which shall consist of the following:

- Date of administration of the vaccine
 - Vaccine manufacturer and lot number of the vaccine
 - Name and title of person administering the vaccine
 - Address of clinic where vaccine was given
 - The address of the facility where the permanent record will reside (if appropriate)
 - Edition date printed on the appropriate VIS
 - Date the VIS was given to the vaccine recipient, or the parents/legal representative
-
- We also recommend that the vaccine type, dose, site, route of administration, and vaccine expiration date be documented, and any vaccine refusal (if appropriate).

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/vac-admin.pdf>

MDPH Vaccine Administration Record

Vaccine Administration Record – All Ages

Record No. / Insurance No.: _____

Patient Name: _____

Address: _____

Birth Date: _____ Male ___ Female ___

Clinic Name and Address:

Use Reverse Side for Names and Initials of Vaccine Administrators

Vaccine administrator: Provide the patient, parent or legal representative with the most recent copy of the Vaccine Information Statement (VIS), which explains risks and benefits of vaccine, for **each** dose of vaccine given.

Type of Vaccine: Record the generic abbreviation for the type of vaccine given (e.g., DTaP), not the trade name. For combination vaccines, indicate the type (e.g., DTaP-Hib) and all other information for each individual antigen (e.g., in the DTP and Hib sections) comprising the combination. Document all lot numbers for each component.

Vaccine	Type of Vaccine	Date Given M/D/Y	Dose	Route (PO, SC, IM, ID, IN, MP)	Site (RA, LA, RT, LT)	Vaccine		Vaccine Information Statement		Vaccine Admin Initials
						lot #	mfr.	Date on VIS	Date Given	
Hepatitis B (e.g., HepB, HepB-Hib, DTaP-HepB-IPV, HepA-HepB)				IM						
				IM						
				IM						
				IM						
Diphtheria, Tetanus, Pertussis (e.g., DTP, DTaP, DT, DTaP-Hib, DTaP-IPV/Hib, DTaP-HepB-IPV, DTaP-IPV, Td, Tdap)				IM						
				IM						
				IM						
				IM						
				IM						
				IM						
Haemophilus influenzae type b (e.g., Hib, HepB-Hib, DTaP-Hib, DTaP-IPV/Hib, Hib-MenCY)				IM						
				IM						
				IM						
				IM						

<http://www.mass.gov/eohhs/docs/dph/cdc/immunization/record-vaccine-admin-clinic.pdf>

MIIS Reporting Requirements

Legislation passed in June 2010, charging MDPH to establish an immunization registry (M.G.L. c. 111, s.24M)

- **Mandatory reporting of all immunizations administered in MA**

Regulations were promulgated January 2015

- outline information on system access, confidentiality, and requirements for data elements to be reported
- describe a provider's duty to inform patients, and a patient's right to object to data sharing across providers

See MIIS table or www.contactmiis.info for more information

Vaccine Adverse Reactions

Adverse reaction

- Extraneous effect caused by vaccine
- Side effect

Adverse event

- Any medical event following vaccination
- May be true adverse reaction
- May be only coincidental

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/genrec.pdf>

Vaccine Adverse Reactions

Local

- Pain, swelling, redness at site of injection
- Occur within a few hours of injection
- Usually mild and self-limited

Systemic

- Fever, malaise, headache
- Nonspecific
- May be unrelated to vaccine

Severe Allergic (anaphylaxis)

- Due to vaccine or vaccine component
- Rare
- Risk minimized by screening

Reporting of Vaccine Errors and Adverse Events

- **VAERS:** Vaccine Adverse Event Reporting System
 - Report all vaccine adverse events to VAERS at vaers.hhs.gov or (800) 822-7967. VAERS report forms are available at vaers.hhs.gov
- **ISMP:** Institute for Safe Medication Practice
 - Report vaccine administration errors (e.g., wrong route, wrong dose, and wrong age) to the (ISMP) via the Vaccine Error Reporting Program (VERP) website <http://ismp.org>.
 - Vaccine administration errors should also be reported to VAERS (as described above), and **MUST** be reported if they resulted in an adverse event.

Vaccine Injury Compensation Program (VICP)

- Established by National Childhood Vaccine Injury Act (1986)
- “No fault” program
- Covers all routinely recommended childhood vaccines
- Vaccine Injury Table
 - Lists conditions associated with each vaccine
 - <http://www.hrsa.gov/vaccinecompensation/vaccineinjurytable.pdf>


<http://www.cdc.gov/vaccines/pubs/pinkbook/safety.html>

Tips to Increase Immunization Rates

Incorporate measures to improve vaccination rates

- Assess immunization status of all patients in every clinical encounter
 - Avoid missed opportunities
- Strongly recommend vaccines that patients need
 - Speak from personal experience
- Administer needed vaccines or refer to a vaccinating provider and confirm receipt
 - Utilize standing orders
 - Offer vaccine only visits
- Provide information in foreign languages
- Document vaccines received by patients, including entering immunization into immunization registry (MIIS)

Many Providers Underestimate the Value Parents Place on Vaccines



Adapted from Healy et al. Vaccine. 2014;32:579-584.

Benefits of Standing Orders

- Overcome administrative barriers and save time
- Shown to be **effective** in both adults and children¹
 - For children, use of standing orders is associated with a median increase in vaccination coverage of 28%
 - Most effective evidence-based method
- **REDUCES MISSED OPPORTUNITIES**
- Consider implementing standing orders for vaccination, particularly for the adolescent immunization 'bundle'
- 'Presumptive' recommendation in action

IAC model standing orders available at:

<http://www.immunize.org/standing-orders/>

MDPH model standing orders available at:

<http://www.mass.gov/eohhs/gov/departments/dph/programs/id/immunization/model-standing-orders.html>


FREQUENTLY ASKED QUESTIONS

MMR Serology for Healthcare Personnel

- Q: Would you consider a healthcare personnel (HCP) with 2 documented doses of MMR vaccine to be considered immune even if their serology for 1 or more of the antigens comes back negative?
- A: Yes, a HCP with 2 documented doses of MMR vaccine are considered to be immune regardless of the results of a subsequent serologic test for measles, mumps, or rubella. Documented age-appropriate vaccination supersedes the results of subsequent serologic testing.
 - HCP who do not have documentation of MMR vaccination and whose serologic test is interpreted as "indeterminate" or "equivocal" should be considered not immune and should receive 2 doses of MMR. ACIP does not recommend serologic testing after vaccination.
- Source: IAC "Ask the Experts, MMR"
http://www.immunize.org/askexperts/experts_mmr.asp

Hepatitis B and Healthcare Personnel

- Guidance for Pre-exposure serologic testing for Healthcare Personnel with a Documented Hepatitis B Vaccine Series Who Have Not Had Post-vaccination Serologic Testing



CDC Guidance:
<https://www.cdc.gov/mmwr/pdf/rr/rr6210.pdf>

IAC's Hepatitis B and Healthcare Personnel:
<http://www.immunize.org/catg.d/p2109.pdf>

Vaccine Preparation “Nevers”

- Never combine vaccine into a single syringe except when specifically approved by the FDA and packaged for that specific purpose
- Never transfer vaccine from one syringe to another
- Never draw partial doses of vaccine from separate vials to obtain a full dose

<https://www.cdc.gov/vaccines/pubs/pinkbook/vac-admin.html#prep>

RESOURCES

General Recommendations on Immunization

Recommendations of the Advisory Committee
on Immunization Practices (ACIP)



Continuing Education Examination available at <http://www.cdc.gov/mmwr/cme/conted.html>



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

ACIP General Recommendations

- Vaccine administration guidelines
- Contraindication and precautions
- Table 1 Recommended and Minimum Ages and Intervals Between Doses and its footnotes

<http://www.cdc.gov/mmwr/pdf/rr/rr6002.pdf>

TABLE 1. Recommended and minimum ages and intervals between vaccine doses*†

Vaccine and dose number	Recommended age for this dose	Minimum age for this dose	Recommended interval to next dose	Minimum interval to next dose
HepB-1 [§]	Birth	Birth	1–4 months	4 weeks
HepB-2	1–2 months	4 weeks	2–17 months	8 weeks
HepB-3 [¶]	6–18 months	24 weeks	—	—
DTaP-1 [§]	2 months	6 weeks	2 months	4 weeks
DTaP-2	4 months	10 weeks	2 months	4 weeks
DTaP-3	6 months	14 weeks	6–12 months	6 months**††
DTaP-4	15–18 months	12 months	3 years	6 months**
DTaP-5	4–6 years	4 years	—	—
Hib-1 ^{§,§§}	2 months	6 weeks	2 months	4 weeks
Hib-2	4 months	10 weeks	2 months	4 weeks
Hib-3 ^{¶¶}	6 months	14 weeks	6–9 months	8 weeks
Hib-4	12–15 months	12 months	—	—
IPV-1 [§]	2 months	6 weeks	2 months	4 weeks
IPV-2	4 months	10 weeks	2–14 months	4 weeks
IPV-3	6–18 months	14 weeks	3–5 years	6 months
IPV-4 ^{***}	4–6 years	4 years	—	—
PCV-1 ^{§§}	2 months	6 weeks	8 weeks	4 weeks
PCV-2	4 months	10 weeks	8 weeks	4 weeks
PCV-3	6 months	14 weeks	6 months	8 weeks

<http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/A/age-interval-table.pdf>

Immunization Action Coalition

www.immunize.org

Home | About IAC | Contact | A-Z Index | Donate | Shop | [SUBSCRIBE](#)

For Health Professionals | For the Public | For Coalitions

Immunization Action Coalition

Google™ Site Search

Handouts for Patients & Staff | Clinic Resources | Vaccine Information Statements | Diseases & Vaccines | Talking about Vaccines | Topics

Welcome

Needle Tips & More

What's New at IAC


Immunization News

Featured Resources


[Shop IAC](#)

Shop IAC

Training video, record cards, and more



New 2012 Laminated Schedules




[Shop IAC](#)

[Get Email Updates](#)

[Make a Donation](#)

[Shop IAC](#)

[Video of the Week](#)



AAP's Risk Communication


Most Popular

WEB SECTIONS

DOWNLOADS

1. Vaccine Information Statements
2. Ask the Experts
3. Handouts for Patients and Staff
4. Photos
5. CDC Schedules
6. IAC Express
7. Shop IAC
8. Clinic Resources
9. Unprotected People Reports
10. Needle Tips
11. Journal Articles
12. Directory of Resources
13. Talking About Vaccines
14. State Laws
15. ACIP Recommendations

IAC Publications

- [Needle Tips](#) **NEW**
 - [Vaccinate Adults](#) **NEW**
 - [IAC Express - Email news](#)
- 


Ask the Experts

Experts from CDC answer challenging and timely questions about vaccines and their administration

- [Questions & Answers](#)
- 

Unprotected People Reports

Real-life accounts of people who have suffered or died from vaccine-preventable diseases: compelling personal testimonies, case reports, and articles



- [Read Reports](#)

Series: Understanding risk communication theory and having tools such as the CASE model encourages fruitful discussion with families about their vaccine safety concerns. The series of videos introduce risk communication and the CASE model, role play two examples of the CASE model in action, and provide feedback on each of the scenarios. These can be viewed individually or as part of a larger group for discussion.

[Visit the VOTW archive](#)

OFFICIAL INFORMATION

[AAP Policy Statements](#)

[ACIP Recommendations](#)

[FDA Product Approval](#)


[State Information](#)

[>> view all](#)

RESOURCES

National Adult Immunization Plan

- Goal 1: Strengthen the adult immunization infrastructure
- Goal 2: Improve access to adult vaccines
- Goal 3: Increase community demand for adult immunizations
- Goal 4: Foster innovation in adult vaccine development and vaccination-related techniques



Learn more at <http://www.hhs.gov/nvpo/national-adult-immunization-plan/index.html>

MA Adult Immunization Coalition(MAIC)

- MAIC is a collaborative partnership dedicated to increasing adult immunization through education, networking, and sharing innovative and best practices.
- There are currently over 200 members representing:
 - Local and state public health organizations
 - Community health centers
 - Health insurance plans
 - Pharmacies
 - Physicians
 - Vaccine manufacturers
 - Long-term-care and senior service organizations
 - Consumer advocacy groups
 - Hospitals
 - Home health
 - College health services



Learn more at
<http://maic.jsi.com/>

Every Child By Two's *State of the ImmUnion*



Regional Immunization Nurses

- **Denise Dillon** – Northeast 978-851-7261
denise.dillon@state.ma.us
- **Linda Jacobs** – Southeast 508-441-3980
linda.jacobs@state.ma.us
- **Lila Coverstone** – Metro Boston & Central Region 617-983-6811
lila.coverstone@state.ma.us
- **Theodora Wohler** – Metro West & Western Region 617-983-6837
theodora.wohler@state.ma.us
- **Katie Reilly**, Nurse Manager 617-983-6833_(T/Th) 508-441-3982_(M/W/F)
catherine.reilly@state.ma.us

MDPH Immunization Program Contact Information

Immunization Program Main Number

For questions about immunization recommendations, disease reporting, etc.

Phone: 617-983-6800

Fax: 617-983-6840

Website: www.mass.gov/dph/imm

MIIS Help Desk

Phone: 617-983-4335

Fax: 617-983-4301

Email: miishelpdesk@state.ma.us

Website: www.contactmiis.info | www.mass.gov/dph/miis

MDPH Vaccine Unit

Phone: 617-983-6828

Fax: 617-983-6924

Email: dph-vaccine-management@state.ma.us

Website: www.mass.gov/dph/imm



**THANK
YOU!**

